

Figure 1.




Figure 2.




Figure 3





Figure 4.


Control.

Row
A



Compound B.

Row
B



1 uM

16 uM

50 uM

100 uM

500 uM





Figure 6.

HPLC profile of soybean cells treated with ^{14}C -carbaryl aspartate alone (control) for 48 hours.



HPLC profile of soybean cells treated with ^{14}C -carbaryl aspartate plus 100 μM compound B for 48 hours.





Figure 7.

Percent ^{14}C -carbamoyl aspartate incorporated
into uracil/uridine.



Percent ^{14}C -carbamoyl aspartate incorporated
into orotate




Figure 9.

Conversion of ¹⁴C-orotate to UMP 30 MAT in corn shoot tissue
with increasing concentrations of PRPP




Figure 10.





Figure 11.

Inhibition of the conversion of 14C-orotate to UMP
with increasing concentrations of pyrazole
aldehyde.



Inhibition of the conversion of 14C-orotate to UMP
with increasing concentrations of barbiturate.




Figure 12.





Figure 13.

Comparing amount of ^{14}C -DHO conversion to orotate 30 MAT in fresh soybean microsomal prep.



Comparing amount of ^{14}C -DHO converted to carbamoyl aspartate between treatments 30 MAT.




Figure 14.




Figure 15.





Figure 16.



<u>R =</u>	<u>Compound</u>	<u>Active and Reversed</u>
	<input type="text" value="F"/>	<input type="text" value="YES"/>
	<input type="text" value="E"/>	<input type="text" value="YES"/>
	<input type="text" value="D"/>	<input type="text" value="YES"/>
	<input type="text" value="J"/>	<input type="text" value="NO"/>




	Scientific Name	Common Name	Bayer Code
Broadleaf weeds	Abutilon theophrasti	velvetleaf	ABUTH
	Ambrosia artemisiifolia	common ragweed	AMBEL
	Galium aparine	catchweed bedstraw	GALAP
	Sesbania exaltata	hemp sesbania	SEBEX
	Solanum nigrum	black nightshade	SOLNI
Grass weeds	Avena fatua	wild oats	AVEFA
	Bromus tectorum	downy brome	BROTE
	Digitaria sanguinalis	large crabgrass	DIGSA
	Echinochloa crus-galli	barnyardgrass	ECHCG
	Setaria viridis	green foxtail	SETVI
	Sorghum halapense (seedling)	johnsongrass	SORHAS
Crops	Glycine max	soybean, Williams var.	GLXMAW
	Oryza sativa	rice, Tebonnet var.	ORYSAT
	Triticum aestivum	winter wheat, Riband var.	TRZAWR
	Zea mays	field corn	ZEAMX




Table 2.



Concentration of the herbicide (μM)



Treatment	500	250	125	63	31	16	7.8	3.9
Compound B	8	7C	6C	6C	5C	5C	5C	5C
Compound B + 100 μM Uracil	5C	3C	3C	0	0	0	0	0
Compound B + 100 μM Uridine	6C	5C	3C	1	0	0	0	0
Compound B + 100 μM UMP	6C	5C	3C	1	0	0	0	0
Compound B + 100 μM AMP	8	7C	6C	5C	5C	5C	5C	5C
Compound B + 100 μM Adenine	8	7C	6C	5C	5C	5C	5C	5C
Compound B + 100 μM Cytosine	8	7C	6C	5C	5C	5C	5C	5C
Compound B + 100 μM Guanine	8	7C	6C	6C	5C	5C	5C	5C
Compound B + 100 μM Thymine	8	7C	6C	6C	5C	5C	5C	5C
Compound B + 100 μM Xanthine	8	7C	6C	6C	5C	5C	5C	5C

Table 3
Percent Inhibition

Compound	structure	% inhibition			
		GH SETVI injury (1 kg/ha post)	Miniscreen injury (re- versal)	Hydroponic xylem in- jury.	uracil re- versal
control		0	0	0	
barbiturate		NT	0	NT	
B		70	60 (0)	70	YES
C		60	80 (50)	75	PARTIAL




Compound	structure	% inhibition			
		GH SETVI injury (1 kg/ha post)	Miniscreen injury (re- versal)	Hydroponic xylem in- jury.	uracil re- versal
D		50	40 (0)	60	YES
E		50	65 (0)	25	YES
F		50	55 (0)	65	YES




Compound	structure	% inhibition			
		GH SETVI injury (1 kg/ha post)	Miniscreen injury (re- versal)	Hydroponic xylem in- jury.	uracil re- versal
G		0	0	20	NO
H		0	0	0	NO




Compound	structure	% inhibition			
		GH SETVI injury (1 kg/ha post)	Miniscreen injury (re- versal)	Hydroponic xylem in- jury.	uracil re- versal
I		0	0	0	NO
J		0	0	0	NO



Notes: NA denotes that compound was not tested.
 Miniscreen data in parenthesis is the extent of uracil reversal at 50 uM tested compound.




Table 4
Percent Inhibition




Compound	structure	% inhibition GH activity on SETVI	% inhibition @50 μ M Miniscreen activity	uracil re- versal
A		70	70	YES
K		60	80	YES
L		60	80	YES

Compound	structure	% inhibition GH activity on SETVI	% inhibition @50 μ M Miniscreen activity	uracil re- versal
M		50	70	YES
N		60	80	YES
O		10	70	YES

Compound	structure	% inhibition GH activity on SETVI	% inhibition @50 μ M Miniscreen activity	uracil re- versal
P		30	80	YES
Q		40	80	YES
R		40	70	YES

Compound	structure	% inhibition GH activity on SETVI	% inhibition @50 μ M Miniscreen activity	uracil re- versal
S		50	80	YES
T		20	70	YES

Compound	structure	% inhibition GH activity on SETVI	% inhibition @50 μ M Miniscreen activity	uracil re- versal
U		0	80	YES
V		50	60	YES
W		40	60	YES

Compound	structure	% inhibition GH activity on SETVI	% inhibition @50 μ M Miniscreen activity	uracil re- versal
X		70	50	YES
Y		50	70	YES
Z		0	30	YES